

METRICS INDUCED BY CERTAIN HILBERT-SCHMIDT FIDELITIES ON POSITIVE SEMI-DEFINITE MATRICES

HO MINH TOAN AND VU THE KHOI

ABSTRACT. Motivated by measuring the *degree of similarity* of a pair of quantum states (density matrices), we consider the metric property of the modified Bures angles and modified Bures distances of symmetric functions which are extensions of some fidelity measures on the spaces \mathcal{P} of nonzero positive semi-definite matrices. We use the positive semi-definiteness of the Gram-type matrices to characterize the metric property of the modified Bures angles. As a consequence, we can show that the modified Bures angles induced by the geometric mean, harmonic mean, minimum and maximum of two positive numbers are metrics on \mathcal{P} . In addition, we can also show that the metric property of the modified Bures angles is stronger than that of the modified Bures distances.

1. INTRODUCTION

A fidelity measure is widely used as a measure of the *degree of similarity* of a pair of quantum states (density matrices). Recently, fidelity measure has been applied to study entanglement quantification, etc. A nice survey of this topic can be found in [1, 2] and the references therein.

If \mathcal{F} is a fidelity measure, we may expect that a metric can be constructed via some functionals of \mathcal{F} . Moreover, we may also expect to study the metric property not only on the quantum state space (the set of density matrices), but also on the space of positive semi-definite matrices Bhatia et al. [6]. The metric property of the modified Bures angle (denoted by $\arccos \sqrt{\mathcal{F}}$), of the modified Bures distance ($\sqrt{1 - \sqrt{\mathcal{F}}}$) and of the modified Bures sine distance ($\sqrt{1 - \mathcal{F}}$) are studied and summarized in [1, Table 6]. Motivated by studying metric properties for some functionals which are considered but still undetermined in [1], we consider that problem for a class of real-valued symmetric functions defined on $\mathcal{P} \times \mathcal{P}$, where

2020 *Mathematics Subject Classification.* 15A45, 15B48, 81P47 .

Key words and phrases. Fidelity, Metric, Bures distance.

\mathcal{P} is the space of nonzero positive semi-definite matrices. Such the functions may be extensions of some fidelity measures on the quantum state spaces. This class includes the well-known fidelity measures such as $\mathcal{F}_{\text{II}}, \mathcal{F}_{\text{AM}}, \mathcal{F}_{\text{GM}}$, etc. (see [1]). Precisely, given a real-valued, symmetric and nonnegative function f on $(\mathbb{R}^+)^2$, we consider the function \mathcal{F}_f defined on $\mathcal{P} \times \mathcal{P}$ as below:

$$(1.1) \quad \mathcal{F}_f(A, B) = \frac{\text{Tr}(AB)}{\sqrt{\text{Tr}(A^2)\text{Tr}(B^2)}} f(\text{Tr}(A^2), \text{Tr}(B^2)),$$

for every pair of nonzero positive semi-definite matrices of the same size A, B . These fidelities measures are called *Hilbert-Schmidt fidelities*, see [1, 2.6]

When we consider metric property of the functions induced by fidelities, the difficult step is to check the triangle inequality. Therefore, we pay attention to the triangle inequality of the modified Bures angles $\arccos \mathcal{F}$ and of the modified Bures distance $\sqrt{1 - \mathcal{F}}$, where \mathcal{F} is defined on \mathcal{P} . In this work, we consider the metric property of the modified Bures angle $\arccos \mathcal{F}_f$. We will show that the modified Bures angle $\arccos \mathcal{F}(A, B)$ satisfies the triangle inequality on \mathcal{P} if and only if the Gram-type matrix of \mathcal{F} is positive semi-definite on \mathcal{P} . As a consequence, if f is the ratio of the geometric mean and maximum mean of two positive numbers then the modified Bures angle $\arccos \mathcal{F}_f$ is a metric on \mathcal{P} . Another application of the above equivalent conditions is to prove the triangle inequality of $\arccos \mathcal{F}_{\text{GM}}$ on \mathcal{P} . Precisely, if f is the constant 1 then $\mathcal{F}_1 = \mathcal{F}_{\text{GM}}$ and the modified Bures angle $\arccos \mathcal{F}_{\text{GM}}$ is a metric on the quantum state space, *not* a metric on \mathcal{P} , even it satisfies the triangle inequality on \mathcal{P} , since $\mathcal{F}_{\text{GM}}(A, B) = 1$ if and only if $\{A, B\}$ is linearly dependent. In other cases, if $f = \frac{m}{\nabla}$, where ∇ is the arithmetic mean of two positive numbers and $m(x, y)$ is a mean which is a member of $\{\min\{x, y\}, \sqrt{xy}, x!y\}$ then we can show that $\arccos \mathcal{F}_f$ is a metric on \mathcal{P} . In addition, we also show that if the modified Bures angle $\arccos \mathcal{F}$ is a metric, so is $\sqrt{1 - \mathcal{F}}$.

The paper is organized as follow: In Section 2, we recall some notations and results obtained in Liang at el. [1]. Section 3 includes the characterization of metric property of the modified Bures angle $\arccos \mathcal{F}_f$ and its applications to some nice cases where f are ratios of means of two positive numbers.

2. PRELIMINARIES AND NOTATIONS

Notations. Let \mathbb{R}^+ denote the set of positive real numbers, M the algebra of $n \times n$ matrices with complex coefficients, \mathcal{P} the space of nonzero positive semi-definite matrices.

The function $\arccos(\mathcal{F})$ ($\sqrt{1 - \mathcal{F}}$) is called the *modified Bures angle* of $\sqrt{\mathcal{F}}$ (the *modified Bures distance* of $\sqrt{\mathcal{F}}$, respectively), see Liang et al. [1].

A large class of functions which are generalizations of the fidelity measures above can be defined as follows. Let f be an arbitrary symmetric, non-vanishing function on $\mathbb{R}^+ \times \mathbb{R}^+$. The function \mathcal{F}_f defined by (1.1) includes some well-known fidelities (these fidelities functions can be also defined on $\mathcal{P} \times \mathcal{P}$) as follows.

Remark. Let \mathcal{F}_f be a function defined by (1.1).

- If $f(x, y) = 2\frac{\sqrt{xy}}{x+y}$, then

$$(2.1) \quad \mathcal{F}_f(A, B) = \frac{2\text{Tr}(AB)}{\text{Tr}(A^2) + \text{Tr}(B^2)} = \mathcal{F}_{\text{AM}}(A, B).$$

- If $f(x, y) = 1$, a constant function then

$$(2.2) \quad \mathcal{F}_f(A, B) = \frac{\text{Tr}(AB)}{\sqrt{\text{Tr}(A^2)}\sqrt{\text{Tr}(B^2)}} = \mathcal{F}_{\text{GM}}(A, B).$$

- If $f(x, y) = \frac{\sqrt{xy}}{\max\{x, y\}}$, then

$$(2.3) \quad \mathcal{F}_f(A, B) = \frac{\text{Tr}(AB)}{\max\{\text{Tr}(A^2), \text{Tr}(B^2)\}} = \mathcal{F}_{\text{II}}(A, B).$$

It is clear that

$$\mathcal{F}_f(A, B) = \mathcal{F}_{\text{GM}}(A, B)f(\text{Tr}(A^2), \text{Tr}(B^2)).$$

Note that the notation \mathcal{F}_f in [1] means the function $\mathcal{F}_{\frac{\#}{f}}$ in the sense (1.1), where $x\#y = \sqrt{xy}$.

We can recall here the results about metric property on the quantum state space for the functionals of the fidelity as in [1, Table 6]:

- Neither $\arccos \sqrt{\mathcal{F}}$ nor $\sqrt{1 - \sqrt{\mathcal{F}}}$ are metrics on the quantum state space if \mathcal{F} is a member in $\{\mathcal{F}_{\text{II}}, \mathcal{F}_{\text{AM}}, \mathcal{F}_{\text{GM}}\}$.
- The function $\sqrt{1 - \mathcal{F}}$ is a metric if \mathcal{F} is a member of $\{\mathcal{F}_{\text{II}}, \mathcal{F}_{\text{GM}}\}$ and is still undetermined if $\mathcal{F} = \mathcal{F}_{\text{AM}}$.

In this paper, we consider metric properties of the functions \mathcal{F}_f on the space of nonzero positive semi-definite matrices \mathcal{P} and similarly we can also obtain such the results on any subspace of \mathcal{P} .

3. RESULTS

Fidelity were introduced in Jozsa [3] and is a mathematical prescription for the quantification of the degree of similarity of a pair of quantum states. The metric property of fidelity measures is an important problem in the study of fidelity. Motivated by this problem in quantum theory, in this note, we study the metric property of the functions \mathcal{F} defined on $\mathcal{P} \times \mathcal{P}$ which should satisfy the *Jozsa's axioms*.

Definition 3.1. Let X be a nonempty set. A real-valued non-vanishing function F defined on $X \times X$ is said to satisfy the *Jozsa's axioms* on X if for every x, y in X , the following conditions hold:

J1a) $0 \leq F(x, y) \leq 1$.

J1b) $F(x, y) = 1$ if and only if $x = y$.

J2) $F(x, y) = F(y, x)$.

Note that the list of Jozsa's axioms in [1] includes some more axioms. However, in this work, we only pay attention to the ones which are useful to the metric property.

3.1. Metric property of the modified Bures angle. We first recall a property from [9, Proposition 1]:

Corollary 3.2. *Let \mathcal{F} be a function defined on $\mathcal{P} \times \mathcal{P}$, which satisfies the Jozsa's axioms **J1a)**, **J1b)** **J2)**. If the modified Bures angle $\arccos \mathcal{F}$ is a metric on \mathcal{P} , so is $\sqrt{1 - \mathcal{F}}$.*

Remark. Let F be a real valued function which satisfies the *Jozsa's axioms* on X . Then the *modified Bures angle* $\arccos F$ satisfies the first two axioms of a metric, i.e.,

- (i) $\arccos F(x, y) \geq 0$ for every $x, y \in X$ and $\arccos F(x, y) = 0$ if and only if $x = y$.
- (ii) $\arccos F(x, y) = \arccos F(y, x)$ for every $x, y \in X$.

Thanks to this remark, to study the metric property of $\arccos F(x, y)$ on X , we only consider the triangle inequality of $\arccos F(x, y)$, i.e.,

$$\arccos F(x, y) \leq \arccos F(x, z) + \arccos F(z, y), \quad \forall x, y, z \in X.$$

Let \mathcal{F} and f be a real valued functions on $\mathcal{P} \times \mathcal{P}$ and $\mathbb{R}^+ \times \mathbb{R}^+$, respectively. $G(\mathcal{F}) = G(\mathcal{F})(A, B, C)$ and $G_f = G_f(x, y, z)$ denote the so-called *Gram-type matrices* of \mathcal{F} and f , respectively and are defined as follows:

$$G(\mathcal{F}) = \begin{pmatrix} \mathcal{F}(A, A) & \mathcal{F}(A, B) & \mathcal{F}(A, C) \\ \mathcal{F}(B, A) & \mathcal{F}(B, B) & \mathcal{F}(B, C) \\ \mathcal{F}(C, A) & \mathcal{F}(C, B) & \mathcal{F}(C, C) \end{pmatrix}, \quad G_f = \begin{pmatrix} f(x, x) & f(x, y) & f(x, z) \\ f(y, x) & f(y, y) & f(y, z) \\ f(z, x) & f(z, y) & f(z, z) \end{pmatrix}$$

for matrices A, B, C in \mathcal{P} and for real numbers x, y, z in \mathbb{R}^+ . Set

$$r(\mathcal{F}) := \{(\mathcal{F}(A, B), \mathcal{F}(B, C), \mathcal{F}(C, A)) \mid A, B, C \in \mathcal{P}\} \subset \mathbb{R}^3.$$

Proposition 3.3. *Let \mathcal{F} be a function which satisfies the Jozsa's axioms **J1a**), **J1b**), **J2**) on \mathcal{P} . Then the following statements are equivalent.*

- (i) $d_{\mathcal{F}} = \arccos(\mathcal{F})$ is a metric on \mathcal{P} .
- (ii) The Gram-type matrix $G(\mathcal{F})$ of \mathcal{F} is positive semi-definite for every positive semidefinite matrices A, B, C in \mathcal{P} .
- (iii)

$$S(x, y, z) := x^2 + y^2 + z^2 - 2xyz - 1 \leq 0, \quad \forall (x, y, z) \in r(\mathcal{F}).$$

Proof. The equivalence of (ii) and (iii) follows from the fact that $S(x, y, z) = \det G(\mathcal{F})$ and the hypothesis that \mathcal{F} satisfies the axiom **J1a**).

(i) \Leftrightarrow (iii). See [9, Proposition 2]. □

Corollary 3.4. *Given a function \mathcal{F} satisfying the Jozsa's axioms **J1a**), **J1b**), **J2**) on \mathcal{P} , if the modified Bures angle $\arccos \mathcal{F}$ is a metric on \mathcal{P} , so is $\arccos(\mathcal{F}^m)$ for every positive integer m .*

Proof. The Gram-type matrix $G(\mathcal{F}^m)$ of \mathcal{F}^m can be decomposed as Schur product of $G(\mathcal{F})$:

$$G(\mathcal{F}^2) = G(\mathcal{F}) \circ G(\mathcal{F}), \quad G(\mathcal{F}^3) = G(\mathcal{F}^2) \circ G(\mathcal{F}) \circ G(\mathcal{F}), \dots$$

Hence, by Schur Product Theorem, $G(\mathcal{F}^m)$ is positive semi-definite. Then the statement follows from Proposition 3.3. □

By the Cauchy Schwarz inequality (with the Frobenius inner product), we have

$$|\mathrm{Tr}(AB)| \leq \sqrt{\mathrm{Tr}(A^2)}\sqrt{\mathrm{Tr}(B^2)}$$

for every A, B in \mathcal{P} . Hence, $0 \leq \mathcal{F}_{\mathrm{GM}}(A, B) \leq 1$. Therefore, the function $\arccos \mathcal{F}_{\mathrm{GM}}(A, B)$ is well-defined. The equality of the Cauchy Schwarz inequality above holds if and only if $A = \lambda B$ for some positive number λ . If we assume further that A, B are density matrices, then $\mathcal{F}_{\mathrm{GM}}(A, B) = 1$ if and only if $A = B$.

Lemma 3.5. *Let $\mathcal{F}_{\mathrm{GM}}$ be the function defined by the formula (2.2). Then the Gram-type matrix of $\mathcal{F}_{\mathrm{GM}}$ is positive semi-definite on \mathcal{P} . As a consequence, the modified Bures angle $\arccos \mathcal{F}_{\mathrm{GM}}$ satisfies the triangle inequality on \mathcal{P} .*

The fidelity $\mathcal{F}_{\mathrm{GM}}$ satisfies the axiom **J1b)** on the space of density matrices but not on \mathcal{P} . Hence, according to the above lemma, $\arccos \mathcal{F}_{\mathrm{GM}}$ is a metric on the quantum state space, but not on \mathcal{P} . Note also that the modified Bures angle $\arccos \sqrt{\mathcal{F}_{\mathrm{GM}}}$ is *not* a metric (it does not satisfy the triangle inequality) on the quantum state space, see [1, Table 6].

Proof. For any nonzero positive semi-definite matrices of the same size A, B, C and let $v(A) = \frac{A}{\sqrt{\mathrm{Tr}(A^2)}}$, $v(B) = \frac{B}{\sqrt{\mathrm{Tr}(B^2)}}$ and $v(C) = \frac{C}{\sqrt{\mathrm{Tr}(C^2)}}$ be the unit vectors in the space M_n with the Frobenius inner product $\langle A, B \rangle = \mathrm{Tr}(B^*A)$. Then the Gram-type matrix of $\mathcal{F}_{\mathrm{GM}}$ is

$$G(\mathcal{F}_{\mathrm{GM}}) = \begin{pmatrix} \langle v(A), v(A) \rangle & \langle v(A), v(B) \rangle & \langle v(A), v(C) \rangle \\ \langle v(B), v(A) \rangle & \langle v(B), v(B) \rangle & \langle v(B), v(C) \rangle \\ \langle v(C), v(A) \rangle & \langle v(C), v(B) \rangle & \langle v(C), v(C) \rangle \end{pmatrix}.$$

Hence, $G(\mathcal{F}_{\mathrm{GM}})$ is positive semi-definite. As in the proof of Proposition 3.3, $\arccos \mathcal{F}_{\mathrm{GM}}$ satisfies the triangle inequality on \mathcal{P} . \square

Theorem 3.6. *Let f be a continuous function which satisfies the Jozsa's axioms **J1a)**, **I1b)** and **J2)** on $\mathbb{R}^+ \times \mathbb{R}^+$. Then the following statements hold.*

- (i) *The function $\mathcal{F}_f(A, B)$ defined by (1.1) satisfies the Jozsas axioms **J1a)**, **J1b)**, **J2)** on \mathcal{P} .*

- (ii) *The modified Bures angle $\arccos \mathcal{F}_f$ is a metric on \mathcal{P} provided that the Gram-type matrix*

$$G_f = \begin{pmatrix} 1 & f(x, y) & f(x, z) \\ f(x, y) & 1 & f(y, z) \\ f(x, z) & f(y, z) & 1 \end{pmatrix}$$

is positive semi-definite for every positive numbers x, y, z .

Proof. (i).

J1a) Combine (1.1), $\mathcal{F}_{\text{GM}}(A, B) \in [0, 1]$ and $f(x, y) \in [0, 1]$, we get $\mathcal{F}_f(A, B) \in [0, 1]$.

J1b) Since $\mathcal{F}_{\text{GM}}(A, B) \leq 1$, $f(x, y) \leq 1$ and the equality (1.1), we have

$$\mathcal{F}_f(A, B) = 1 \Leftrightarrow \mathcal{F}_{\text{GM}}(A, B) = 1 \text{ and } f(\text{Tr}(A^2), \text{Tr}(B^2)) = 1.$$

Since $f(x, y) = 1$ if and only if $x = y$, we get that $\text{Tr}(A^2) = \text{Tr}(B^2)$. In addition, $\mathcal{F}_{\text{GM}}(A, B) = 1$ if and only if $A = cB$ for some constant $c > 0$. Hence, A must be equal to B .

J2) The symmetry of \mathcal{F}_f follows from that of \mathcal{F}_{GM} and of f .

(ii) By Proposition 3.3, the modified Bures angle $\arccos \mathcal{F}_f$ is a metric if and only if the Gram-type matrix $G(\mathcal{F}_f)$ is positive semi-definite. Since

$$\mathcal{F}_f(A, B) = \mathcal{F}_{\text{GM}}(A, B)f(\text{Tr}(A^2), \text{Tr}(B^2)),$$

we can write $G(\mathcal{F}_f)$ as the Schur product:

$$G(\mathcal{F}_f) = G(\mathcal{F}_{\text{GM}}) \circ G_f.$$

The matrix $G(\mathcal{F}_{\text{GM}})$ is positive semi-definite by Lemma 3.5, while G_f is positive semi-definite by the hypothesis. These together with Schur's Product Theorem imply that $G(\mathcal{F}_f)$ is positive semi-definite. \square

Note: In Theorem 3.6, $\arccos \mathcal{F}_f$ is a metric on *the quantum state space* even when we remove the condition ' $f(x, y) = 1 \Rightarrow x = y$ ' in the hypothesis of the theorem.

Example. If $f(x, y)$ is a real-valued positive definite kernel, then the Gram-type matrix G_f is positive semi-definite. In particular, if $g(x)$ is an even nonnegative function on \mathbb{R} then $f(x, y) = \exp(-g(x - y))$ is a positive definite kernel. Furthermore, $f(x, y)$ satisfies the Jozsa's axioms on \mathbb{R}^+ provided that 0 is the unique zero

point of $g(x)$. By Theorem 3.6, $\arccos \mathcal{F}_f$ is a metric on \mathcal{P} and so is $\sqrt{1 - \mathcal{F}_f}$. We may find more interesting examples of positive definite functions/kernels in [7, 8].

Corollary 3.7. *The modified Bures angle $\arccos \mathcal{F}_{II}$ is a metric on \mathcal{P} , where \mathcal{F}_{II} is defined by (2.3).*

Note that neither $\arccos \sqrt{\mathcal{F}_{II}}$ nor $\sqrt{1 - \sqrt{\mathcal{F}_{II}}}$ are metrics, see [1, Table 6].

Proof. Let $f(x, y) = \frac{\sqrt{xy}}{\max\{x, y\}}$. Then f satisfies the Jozsa's axioms on $\mathbb{R}^+ \times \mathbb{R}^+$ and

$$\mathcal{F}_f(A, B) = \mathcal{F}_{II}(A, B).$$

Let's consider the Gram-type matrix

$$G_f = \begin{pmatrix} 1 & f(x, y) & f(x, z) \\ f(y, x) & 1 & f(y, z) \\ f(z, x) & f(z, y) & 1 \end{pmatrix},$$

Denote by $a = f(x, y)$, $b = f(y, z)$ and $c = f(z, x)$. Without loss of generality, we can assume that $0 \leq x \leq y \leq z \leq 1$, we have

$$a = f(x, y) = \frac{\sqrt{x}}{\sqrt{y}}, \quad b = f(y, z) = \frac{\sqrt{y}}{\sqrt{z}}, \quad c = f(z, x) = \frac{\sqrt{x}}{\sqrt{z}}.$$

Then

$$\det G_f = 1 + 2abc - a^2 - b^2 - c^2 = 1 + \frac{x}{z} - \frac{x}{y} - \frac{y}{z} \geq 0,$$

since $\frac{x}{y} \leq \frac{x+z-y}{z}$. Therefore, G_f is positive semi-definite. By Theorem 3.6, the modified Bures angle $\arccos \mathcal{F}_{II}$ is a metric on \mathcal{P} . \square

3.2. Fidelity measures induced by means.

3.2.1. *Means of positive numbers.* A mean of positive numbers is a function $m : \mathbb{R}^+ \times \mathbb{R}^+ \rightarrow \mathbb{R}^+$ which satisfies the following conditions:

- (i) $m(x, x) = x$ for every positive number x .
- (ii) $m(x, y) = m(y, x)$ for every positive numbers x, y .
- (iii) If $x < y$ then $x < m(x, y) < y$.
- (iv) If $x < x'$ and $y < y'$ then $m(x, y) < m(x', y')$.
- (v) $m(x, y)$ is continuous.

A mean is said to be *homogeneous* if

- (vi) $m(tx, ty) = tm(x, y)$ for every positive numbers x, y, t .

See [4, 5] for more details. Thanks to the homogeneity, a two variable function m can be reduced to a single variable function m_1 such that

$$m(x, y) = xm_1\left(\frac{y}{x}\right) = ym_1\left(\frac{x}{y}\right).$$

Furthermore, a homogeneous mean m above is uniquely described by such a function m_1 satisfying the following properties:

- (i)' $m_1(1) = 1$.
- (ii)' $tm_1(t^{-1}) = m_1(t)$.
- (iii)' $m_1(t) > 1$ if $t > 1$ and $m_1(t) < 1$ if $0 < t < 1$.
- (iv)' m_1 is monotone increasing.
- (v)' m_1 is continuous.

Conversely, each function f satisfying the properties above gives a mean $m(x, y) = xm_1\left(\frac{y}{x}\right)$.

The following means are the most well-known ones:

- (1) The geometric mean $x\sharp y := \sqrt{xy}$ whose corresponding function \sqrt{t} .
- (2) The arithmetic mean $x\nabla y := (x+y)/2$ whose corresponding function $(t+1)/2$.
- (3) The harmonic mean $x!y := 2(x^{-1} + y^{-1})^{-1}$ whose corresponding function $2t/(1+t)$.

Proposition 3.8. *Let $m \leq \sharp$ be a homogeneous mean (of two positive numbers). Denote by $\mathcal{F}_{\frac{m}{\nabla}}$ the function defined by the formula (1.1).*

- (i) *The function $\mathcal{F}_{\frac{m}{\nabla}}$ satisfies the Jozsa's axioms on \mathcal{P} .*
- (ii) *The modified Bures distance $\mathcal{F}_{\frac{m}{\nabla}}$ is a metric provided that*

$$(3.1) \quad \frac{1}{4} + 4 \frac{m_1(s)}{1+s} \frac{m_1(t)}{1+t} \frac{m_1(st)}{1+st} \geq \left(\frac{m_1(s)}{1+s}\right)^2 + \left(\frac{m_1(t)}{1+t}\right)^2 + \left(\frac{m_1(st)}{1+st}\right)^2$$

holds true for all positive numbers s, t , where $m_1(x) = m(1, x)$.

Proof. Let $f(x, y) = \frac{2m(x, y)}{x+y}$ be a real-valued function on $\mathbb{R}^+ \times \mathbb{R}^+$. Then $\mathcal{F}_{\frac{m}{\nabla}} = \mathcal{F}_f$ defined by (1.1).

(i) Since $m(x, y) = m(y, x)$, $f(x, y)$ is a nonnegative symmetric function on $(\mathbb{R}^+)^2$. As $m(x, y) \leq x\nabla y$, we have $0 \leq f(x, y) \leq 1$. Clearly $f(x, x) = 1$ because $m(x, x) = x = x\nabla x$. Suppose that $f(x, y) = 1$. Equivalently, $m(x, y) = x\nabla y$. Since $m \leq \sharp$, we have $\sqrt{xy} = x\nabla y$. Hence, $x = y$. Therefore, f satisfies the Jozsa's axioms on \mathbb{R}^+ .

Thus, by Theorem 3.6, $\mathcal{F}_{\frac{m}{\nabla}}$ is a fidelity measure satisfying **J1a)**, **J1b)**, **J2)**.

(ii) By Theorem 3.6, $\arccos \mathcal{F}_{\frac{m}{\nabla}}$ is a metric if the Gram-type matrix of $f = \frac{m}{\nabla}$

$$G_f = \begin{pmatrix} 1 & f(x, y) & f(x, z) \\ f(x, y) & 1 & f(y, z) \\ f(x, z) & f(y, z) & 1 \end{pmatrix}$$

is positive semi-definite. This is equivalent to $\det G_f \geq 0$, i.e.,

$$(3.2) \quad 1 + 2f(x, y)f(y, z)f(z, x) \geq f^2(x, y) + f^2(y, z) + f^2(z, x), \quad \forall x, y, z \in (0, 1].$$

We have

$$f(x, y) = \frac{2m(x, y)}{x + y} = \frac{2m_1(s)}{1 + s}, \quad s = \frac{x}{y}.$$

Similarly, $f(y, z) = \frac{2m_1(t)}{1+t}$ with $t = \frac{y}{z}$ and $f(x, z) = \frac{2m_1(st)}{1+st}$. Substituting these identities into the inequality (3.2), we get (3.1). \square

Corollary 3.9. *The modified Bures angle $\arccos \mathcal{F}_{AM}$ is a metric on \mathcal{P} , where \mathcal{F}_{AM} is defined by (2.1).*

Proof. See the proof of [9, Theorem 4]. \square

Corollary 3.10. *Let $f(x, y)$ be a symmetric function defined on $(\mathbb{R}^+)^2$ and \mathcal{F} by*

$$\mathcal{F}(A, B) = \frac{2\text{Tr}(AB)}{\text{Tr}(A^2) + \text{Tr}(B^2)} f(\text{Tr}A^2, \text{Tr}B^2)$$

for every pair of the same size matrices $A, B \in \mathcal{P}$. Suppose that $0 \leq f(x, y) \leq 1$, $f(x, x) = 1$ for every positive numbers x, y and the Gram-type matrix G_f is positive definite on \mathbb{R}^+ . Then $\arccos \mathcal{F}$ is a metric on \mathcal{P} .

Proof. Since $\mathcal{F}(A, B) = \mathcal{F}_{AM}(A, B)f(\text{Tr}A^2, \text{Tr}B^2)$, and \mathcal{F}_{AM} satisfies the Jozsa's axioms **J1a**), **J1b**) and **J2**), $\arccos \mathcal{F}(A, B)$ is nonnegative, symmetric and is equal to zero if and only if $A = B$. Since $\arccos \mathcal{F}_{AM}$ is a metric on \mathcal{P} and by Theorem 3.6, the Gram-type matrix $G(\mathcal{F}_{AM})$ is positive semi-definite. Thus, $G(\mathcal{F}) = G(\mathcal{F}_{AM}) \circ G_f$ is positive semi-definite by the hypothesis, Corollary 3.9 and Schur product Theorem. Now, apply Theorem 3.6 again, we get the proof. \square

Corollary 3.11. *The modified Bures angle $\arccos \mathcal{F}_{\frac{1}{\nabla}}$ is a metric on \mathcal{P} , where $\frac{1}{\nabla}(x, y) = 2\frac{xy}{x+y}$ is the ratio of the harmonic and arithmetic means.*

Proof. By definition, we have

$$\mathcal{F}_{\frac{1}{\sqrt{\cdot}}}(A, B) = \mathcal{F}_{\text{AM}}(A, B) \frac{x!y}{\sqrt{xy}},$$

where $x = \text{Tr}(A^2), y = \text{Tr}(B^2)$. By Corollary 3.10, it suffices to show that the Gram-type matrix G_f is positive semi-definite, where $f(x, y) = \frac{x!y}{\sqrt{xy}}$. We can write

$$f(x, y) = \frac{x!y}{\sqrt{xy}} = \frac{2}{\sqrt{xy}(x^{-1} + y^{-1})} = \frac{2\sqrt{s}}{1+s}, \quad \text{where } s = \frac{x}{y}.$$

Similarly,

$$f(y, z) = \frac{2\sqrt{t}}{1+t}, \quad f(x, z) = \frac{2\sqrt{st}}{1+st}, \quad \text{where } t = \frac{y}{z}.$$

The fact is that $0 < \min\{x, y\} \leq x!y \leq \sqrt{xy} \leq \max\{x, y\}$, so $0 < f(x, y) \leq 1$. Hence, the Gram-type matrix G_f of f is positive semi-definite (for all positive numbers s, t) if and only if $\det G_f \geq 0$. Equivalently,

$$(3.3) \quad 1 + 2 \frac{2\sqrt{s}}{s+1} \frac{2\sqrt{t}}{t+1} \frac{2\sqrt{st}}{st+1} - 4 \left[\frac{s}{(s+1)^2} + \frac{t}{(t+1)^2} + \frac{st}{(st+1)^2} \right] \geq 0 \quad \text{for } s, t > 0.$$

As the same argument in the proof of Corollary 3.9, the inequality (3.3) holds true for all positive numbers s, t . \square

If we relax the condition (iii) in the definition of mean above by

$$(iii)'' \quad \text{If } x \leq y \text{ then } x \leq m(x, y) \leq y$$

then the function \max, \min are also means of two positive numbers. Actually, Proposition 3.8 still holds true for such a relaxed mean m provided that m can be written as $m(x, y) = xm_1(y/x) = ym_1(x/y)$.

Corollary 3.12. *Let $\mathcal{F} = \mathcal{F}_f$ be a function defined by (1.1). If $f(x, y) = \frac{2\min\{x, y\}}{x+y}$ then $\arccos \mathcal{F}$ is a metric on \mathcal{P} .*

Proof. We can write

$$\mathcal{F}(A, B) = \mathcal{F}_{\text{AM}}(A, B)g(\text{Tr}A^2, \text{Tr}B^2),$$

where $g(x, y) = \frac{\min\{x, y\}}{\sqrt{xy}}$. Let x, y, z be arbitrary positive numbers, without loss of generality, we can assume that $x \leq y \leq z$. Then

$$g(x, y) = \frac{\sqrt{x}}{\sqrt{y}}, \quad g(x, z) = \frac{\sqrt{x}}{\sqrt{z}} \quad \& \quad g(y, z) = \frac{\sqrt{y}}{\sqrt{z}}.$$

Hence the Gram-type matrix G_g is positive semi-definite (as the same argument in the proof of Corollary 3.7). By Corollary 3.10, $\arccos \mathcal{F}$ is a metric on \mathcal{P} . \square

ACKNOWLEDGMENT

The second author is partially supported by Grant number ICRTM02-2021.01 of International Center for Research and Postgraduate Training in Mathematics, HIM, VAST. The authors would like to thank the Vietnam Institute for Advanced Study in Mathematics for the hospitality during the writing of this paper.

REFERENCES

- [1] Y. C. Liang, Y. H. Yeh, P. E. M. F. Mendon, R. Y. Teh, M. D. R. and P. D. Drummond, *Quantum fidelity measures for mixed states*, Rep. Prog. Phys. **82**(7) (2019), 076001.
- [2] S. J. Gu, *Fidelity approach to quantum phase transitions*, Int. J. Mod. Phys. B, **24**(23) (2010), 4371-4458.
- [3] R. Jozsa, *Fidelity for mixed quantum states*, J. Mod. Opt., **41**(12) (1994), 2315- 2323.
- [4] D. Petz and R. Temesi, *Means of positive numbers and matrices*, SIAM J. Matrix Anal. Appl. Vol. **27**, No. 3 (2005), 712-720.
- [5] F. Kubo, T. Ando, *Means of positive linear operators*, Math. Ann. **246** (1980), 205-224.
- [6] R. Bhatia, T. Jain and Y. Lim, *On the Bures-Wasserstein distance between positive definite matrices*, Expo. Math., **37**(2) (2019),165-191.
- [7] R. Bhatia and K. R. Parthasarathy, *Positive definite functions and operator inequalities*, Bull. London Math. Soc. **32**(2) (2000), 214-228.
- [8] T. Hofmann, B. Scholkopf, and A. Smola, *Kernel methods in machine learning*, Ann. Stat., **36**(3) (2008), 1171.
- [9] Vu The Khoi, Ho M. Toan, *Metric properties of alternative fidelities*, Quantum Information and Computation, Vol. 22, No. **9** & **10** (2022) 790-799.

INSTITUTE OF MATHEMATICS (VAST), 18 HOANG QUOC VIET, 10307, HANOI, VIETNAM
E-mail address: hmtoan@math.ac.vn

INSTITUTE OF MATHEMATICS (VAST), 18 HOANG QUOC VIET, 10307, HANOI, VIETNAM
E-mail address: vt Khoi@math.ac.vn